


Doctor Discussion Guide

This is a guide to help you talk with your doctor or healthcare provider about eczema and if EBGLYSS may be right for you.

Before your next doctor's appointment, please take a moment to print or fill out the questions directly in this PDF.

Where do you most often experience your symptoms?
(Check all that apply)



- Face/Scalp
- Neck
- Torso
- Elbows/Arms
- Back
- Buttocks/Genitals
- Hands
- Knees/Legs
- Ankles
- Feet

INDICATION

EBGLYSS is a prescription medicine used to treat adults and children 12 years of age and older who weigh at least 88 pounds (40kg) with moderate-to-severe eczema (atopic dermatitis) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. EBGLYSS can be used with or without topical corticosteroids.

It is not known if EBGLYSS is safe and effective in children less than 12 years of age or in children 12 years to less than 18 years of age who weigh less than 88 pounds (40 kg).

SELECT IMPORTANT SAFETY INFORMATION

Do not use EBGLYSS if you are allergic to lebrikizumab-lbkz or to any of the ingredients in EBGLYSS. See the Patient Information leaflet that comes with EBGLYSS for a complete list of ingredients.

Please see Indication and Safety Summary on page 5.

Doctor Discussion Guide

Over the last week, how many days did you experience eczema symptoms? (Check one)

0 days 1-2 days 3-5 days 6-7 days

Over the last week, how many days did you have intense episodes of itching because of eczema? (Check one)

0 days 1-2 days 3-5 days 6-7 days

What You CAN See

Based on your experience over the last week, select the severity of the following signs of eczema:

Dry/flaky skin

None											Severe
0	1	2	3	4	5	6	7	8	9	10	

Discoloration/redness

None											Severe
0	1	2	3	4	5	6	7	8	9	10	

Scratch marks/bleeding

None											Severe
0	1	2	3	4	5	6	7	8	9	10	

SELECT IMPORTANT SAFETY INFORMATION

Before using EBGLYSS, tell your healthcare provider about all your medical conditions, including if you:

- Have a parasitic (helminth) infection.
- Are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with EBGLYSS.
- Are pregnant or plan to become pregnant. It is not known if EBGLYSS will harm your unborn baby. If you become pregnant during treatment with EBGLYSS, you or your healthcare provider can call Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979) to report the pregnancy.
- Are breastfeeding or plan to breastfeed. It is not known if EBGLYSS passes into your breast milk.
- Take prescription or over-the-counter medicines, vitamins, and herbal supplements.

Please see Indication and Safety Summary on page 5.

Doctor Discussion Guide

What You CAN'T See

Based on your experience over the last week, select the severity of your itching:

Itchiness

No itch											Worst itch imaginable
0	1	2	3	4	5	6	7	8	9	10	

How does eczema impact you?

Treatment Goals

Which treatments have you used in the past? (Check all that apply)

Oral or injected steroids

Phototherapy

Biologic injections

Topical corticosteroids

Over-the-counter products

Other (Please specify):

What do you expect from your next eczema treatment? (Check all that apply)

Clearer skin

More itch relief

Lower my steroid and/or topical use

Long-lasting relief

A dosing schedule that I can maintain

Other (Please specify):

SELECT IMPORTANT SAFETY INFORMATION

EBGLYSS can cause serious side effects, including:

- **Allergic reactions. EBGLYSS can cause allergic reactions that may sometimes be severe.** Stop using EBGLYSS and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms:
 - breathing problems or wheezing
 - swelling of the face, lips, mouth, tongue or throat
 - hives
 - itching
 - fainting, dizziness, feeling lightheaded
 - skin rash
 - cramps in your stomach area (abdomen)
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision.

Please see Indication and Safety Summary on page 5.

Questions for Your Doctor

To prepare for a smooth conversation with your doctor, familiarize yourself with the questions below and select the ones you'd like to ask.

Living with Eczema

What can I do in my daily life to help with my eczema?

How does skin tone change how my eczema should be treated?

Are there any types of eczema treatments that I haven't tried yet?

Understanding EBGLYSS

What is a biologic?

What is IL-13?

Why might EBGLYSS be a good fit for me?

How well does EBGLYSS work, and how long can those results last?

Are there any resources available to help me get started?

Safety & Important Considerations

What side effects might I experience with EBGLYSS?

Can I use EBGLYSS with my other medicines or treatments?

How often should my appointments be while I'm taking EBGLYSS?

Savings & Support

How can I find out if my insurance will cover EBGLYSS?

What support options are available through Lilly Support Services™ that may help with my treatment journey?

PRO TIP: If you are eligible and commercially insured, you may be able to save on EBGLYSS.

Click [here](#) for more information or visit <https://ebglyss.lilly.com/savings-support>

Have a question that wasn't listed above? Write it here so you don't forget it!

SELECT IMPORTANT SAFETY INFORMATION

The most common side effects of EBGLYSS include:

- eye and eyelid inflammation, including redness, swelling, and itching
- injection site reactions
- shingles (herpes zoster)

These are not all of the possible side effects of EBGLYSS. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see Indication and Safety Summary on page 5.

Indication and Safety Summary

EBGLYSS™ (EHB-glihs) is an injectable medicine used to treat adults and children 12 years of age and older who weigh at least 88 pounds (40 kg) with moderate-to-severe eczema (atopic dermatitis) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. EBGLYSS can be used with or without topical corticosteroids.

It is not known if EBGLYSS is safe and effective in children less than 12 years of age or in children 12 years to less than 18 years of age who weigh less than 88 pounds (40 kg).

WARNINGS - Do not use EBGLYSS if you are allergic to lebrikizumab-lbkz or to any of the ingredients in EBGLYSS. See the Patient Information leaflet that comes with EBGLYSS for a complete list of ingredients.

BEFORE USING

Before using EBGLYSS, tell your healthcare provider about all your medical conditions, including if you:

- Have a parasitic (helminth) infection.
- Are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with EBGLYSS.
- Are pregnant or plan to become pregnant. It is not known if EBGLYSS will harm your unborn baby. If you become pregnant during treatment with EBGLYSS, you or your healthcare provider can call Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979) to report the pregnancy.
- Are breastfeeding or plan to breastfeed. It is not known if EBGLYSS passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

POSSIBLE SIDE EFFECTS

EBGLYSS can cause serious side effects, including:

- **Allergic reactions. EBGLYSS can cause allergic reactions that may sometimes be severe.** Stop using EBGLYSS and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms:
 - breathing problems or wheezing
 - swelling of the face, lips, mouth, tongue or throat
 - hives
 - itching
 - fainting, dizziness, feeling lightheaded
 - skin rash
 - cramps in your stomach area (abdomen)
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision.

The most common side effects of EBGLYSS include:

- eye and eyelid inflammation, including redness, swelling, and itching
- injection site reactions
- shingles (herpes zoster)

These are not all of the possible side effects of EBGLYSS.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

HOW TO TAKE

- See the detailed “Instructions for Use” that comes with EBGLYSS for information about how to prepare and inject EBGLYSS and how to properly store and throw away (dispose of) used EBGLYSS prefilled pens and prefilled syringes.
 - Use EBGLYSS exactly as prescribed by your healthcare provider.
 - EBGLYSS is given as an injection under the skin (subcutaneous injection).
 - If your healthcare provider decides that you or a caregiver can give the injections of EBGLYSS, you or a caregiver should receive training on the right way to prepare and inject EBGLYSS. Do not try to inject EBGLYSS until you have been shown the right way by your healthcare provider. In children 12 years of age and older, EBGLYSS should be given by a caregiver.
 - If you miss a dose of EBGLYSS, inject the missed dose as soon as possible, then inject your next dose at your regular scheduled time.

LEARN MORE

EBGLYSS is a prescription medicine available as a 250 mg/2 mL injection prefilled pen or prefilled syringe. For more information, call 1-800-545-5979 or go to ebglyss.lilly.com

This summary provides basic information about EBGLYSS but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking to your doctor. Be sure to talk to your doctor or other healthcare provider about EBGLYSS and how to take it. Your doctor is the best person to help you decide if EBGLYSS is right for you.

LK CON BS AD APP

EBGLYSS™, its delivery device base, and Lilly Support Services™ are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.